

Batteries - Wet Filled with Acid

EC10(ECx)	168h	Algae or other aquatic plants	0.0046mg/l	2
EC50	96h	Algae or other aquatic plants	0.11-0.209mg/l	4
EC50	72h	Algae or other aquatic plants	0.254mg/l	2
LC50	96h	Fish	2.8-4.2mg/l	Not Available

Legend: *Extracted from 1. IUCLID Toxicity Data 2. Europe ECHA Registered Substances - Ecotoxicological Information - Aquatic Toxicity 4. US EPA, Ecotox database - Aquatic Toxicity Data 5. ECETOC Aquatic Hazard Assessment Data 6. NITE (Japan) - Bioconcentration Data 7. METI (Japan) - Bioconcentration Data 8. Vendor Data*

Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment.

Do NOT allow product to come in contact with surface waters or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash-waters.

Wastes resulting from use of the product must be disposed of on site or at approved waste sites.

Ecotoxicity:

The tolerance of water organisms towards pH margin and variation is diverse. Recommended pH values for test species listed in OECD guidelines are between 6.0 and almost 9. Acute testing with fish showed 96h-LC50 at about pH 3.5

Speciation of arsenic is an important consideration in the fate, movement, and action of this substance. Chemical and biochemical transformations of arsenic include oxidation, reduction and methylation which affects its volatilisation, adsorption, dissolution and biological disposition. The transport of arsenic in the environment is largely controlled by absorption/desorption processes in soils and sediments. Sediment movement is responsible for transport of arsenic soil residues to their ultimate sinks in deep ocean sediments. The clay fraction, plus ferrous and aluminium oxides which coat clay particles, adsorb arsenicals which then undergo transformation as discussed earlier. Conversions of arsenic to volatile alkylarsines leads to air transport losses from soil. Inorganic arsenic occurs in water in different oxidation states depending on the pH and Eh of the water. Arsenate is apparently reduced by bacteria to arsenite in marine environments because the ratio of total arsenate to total arsenic is much lower than that predicted thermodynamically. Methylation of arsenic occurs in both freshwater and marine systems where arsenate occurs as arsenate, arsenite, methanearsonic acid and dimethylarsinic acid. Arsenate predominates because this is the most stable form.

Bioaccumulation occurs in some aquatic species such as seaweeds, freshwater algae and crustaceans. Some arsenic in water flea (Daphnia Magna) and algae occurs as arseno-analogues of phospholipids leading to the mistaken impression that accumulation and utilisation of arsenic takes place at the expense of phosphate. Crabs, lobsters and other marine organisms accumulate organo-arsenicals in the food chain. Although human activity may alter the local picture of environmental arsenic there is little evidence that this affects the global scale arsenic cycle.

Airborne concentrations of arsenic in urban areas may range from a few nanograms to a few tenths of a microgram per cubic meter; airborne arsenic is generally inorganic. In oxygenated soils inorganic arsenic is present in pentavalent form; under reducing conditions it occurs as the trivalent form.

Drinking Water Standards:

arsenic: 50 ug/l (UK max.) 0.01mg/L (WHO provisional guideline)

chloride:400 mg/l (UK max.)

250 mg/l (WHO guideline)

Soil Guideline:

Dutch Criteria: 29 mg/kg (target)

55 mg/kg (intervention)

Air Quality Standard:

No safe levels recommended due to carcinogenic properties (WHO guideline)

Lead is primarily an atmospheric pollutant that enters soil and water as fallout, a process determined by physical form and particle size. Lead in the form of alkyls has been introduced to the environment primarily from leaded petrol. These are converted to water-soluble lead compounds of high toxicity and availability to plants. Such compounds easily leach from soil to contaminate water sources close to highways. Lead that has entered the aquatic system from run-off or as fallout of insoluble precipitates is found in sediments. The biological methylation of inorganic lead by lake sediment micro-organisms has been demonstrated although its significance is not entirely clear. Other forms of soluble or insoluble lead may also enter the environment and undergo bioaccumulation through a series of biological incidents.

Sulfuric acid is soluble in water and remains indefinitely in the environment as sulfate.

Large discharges may contribute to the acidification of water and be fatal to aquatic life and soil micro-organisms. Large discharges may also contribute to the acidification of effluent treatment systems and injure sewage treatment organisms.

In water, sulfuric acid dissociates, and the sulfate anion may combine with other cations. In soil, the ions from sulfuric acid can adsorb to soil particles or leach into surface water and groundwater. Sulfates can be taken up by plants and be incorporated into the parenchyma of the plant.

The ions (sulfate, hydrogen) can adsorb to soil particles or be converted to gases. Anaerobic bacteria in sediments and soil can reduce sulfate to sulfur and hydrogen sulfide.

Sulfates, including sulfuric acid, are removed from the air by both dry and wet deposition processes. Wet deposition processes including rain-out (a process that occurs within the clouds) and washout (removal by precipitation below the clouds) contribute to the removal of sulfate from the atmosphere. In the stratosphere, sulfuric acid aerosols have lifetimes of about 14 and 2.4 days at altitudes of 15 and 20 km, respectively. At cloud level, the residence time is about 6 days, with shorter residence times in surface air.

Prevent, by any means available, spillage from entering drains or water courses.

DO NOT discharge into sewer or waterways.

Persistence and degradability

Ingredient	Persistence: Water/Soil	Persistence: Air
	No Data available for all ingredients	No Data available for all ingredients

Bioaccumulative potential

Ingredient	Bioaccumulation

Ingredient	Bioaccumulation
	No Data available for all ingredients

Mobility in soil

Ingredient	Mobility
	No Data available for all ingredients

SECTION 13 Disposal considerations

Waste treatment methods

Product / Packaging disposal	<ul style="list-style-type: none"> ▶ Recycle wherever possible or consult manufacturer for recycling options. ▶ Consult State Land Waste Management Authority for disposal. ▶ DO NOT allow wash water from cleaning or process equipment to enter drains. ▶ It may be necessary to collect all wash water for treatment before disposal. ▶ In all cases disposal to sewer may be subject to local laws and regulations and these should be considered first. ▶ Where in doubt contact the responsible authority. ▶ Recycle wherever possible. ▶ Consult manufacturer for recycling options or consult local or regional waste management authority for disposal if no suitable treatment or disposal facility can be identified. ▶ Treat and neutralise at an approved treatment plant. Treatment should involve: Mixing or slurring in water; Neutralisation followed by: burial in a land-fill specifically licensed to accept chemical and / or pharmaceutical wastes or Incineration in a licensed apparatus (after admixture with suitable combustible material) ▶ Decontaminate empty containers. Observe all label safeguards until containers are cleaned and destroyed.
-------------------------------------	---

SECTION 14 Transport information

Labels Required

	 
Marine Pollutant	NO
HAZCHEM	2R

Land transport (ADG)

UN number or ID number	2794	
UN proper shipping name	BATTERIES, WET, FILLED WITH ACID, electric storage	
Transport hazard class(es)	Class	8
	Subsidiary risk	6.1
Packing group	Not Applicable	
Environmental hazard	Not Applicable	
Special precautions for user	Special provisions	295 AU08
	Limited quantity	1 L

Air transport (ICAO-IATA / DGR)

UN number	2794	
UN proper shipping name	Batteries, wet, filled with acid electric storage	
Transport hazard class(es)	ICAO/IATA Class	8
	ICAO / IATA Subrisk	6.1
	ERG Code	8L
Packing group	Not Applicable	
Environmental hazard	Not Applicable	

Batteries - Wet Filled with Acid

Special precautions for user	Special provisions	A51 A164 A183 A802
	Cargo Only Packing Instructions	870
	Cargo Only Maximum Qty / Pack	400 kg
	Passenger and Cargo Packing Instructions	870
	Passenger and Cargo Maximum Qty / Pack	30 kg
	Passenger and Cargo Limited Quantity Packing Instructions	Forbidden
	Passenger and Cargo Limited Maximum Qty / Pack	Forbidden

Sea transport (IMDG-Code / GGVSee)

UN number	2794	
UN proper shipping name	BATTERIES, WET, FILLED WITH ACID electric storage	
Transport hazard class(es)	IMDG Class	8
	IMDG Subrisk	6.1
Packing group	Not Applicable	
Environmental hazard	Not Applicable	
Special precautions for user	EMS Number	F-A, S-B
	Special provisions	295
	Limited Quantities	1 L

Transport in bulk according to Annex II of MARPOL and the IBC code

Not Applicable

Transport in bulk in accordance with MARPOL Annex V and the IMSBC Code

Product name	Group
lead	Not Available
antimony	Not Available
calcium	Not Available
tin fume	Not Available
sulfuric acid	Not Available
arsenical dust	Not Available

Transport in bulk in accordance with the IGC Code

Product name	Ship Type
lead	Not Available
antimony	Not Available
calcium	Not Available
tin fume	Not Available
sulfuric acid	Not Available
arsenical dust	Not Available

SECTION 15 Regulatory information

Safety, health and environmental regulations / legislation specific for the substance or mixture

lead is found on the following regulatory lists

Batteries - Wet Filled with Acid

Australia Hazardous Chemical Information System (HCIS) - Hazardous Chemicals

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 4

Australian Inventory of Industrial Chemicals (AIIC)

Chemical Footprint Project - Chemicals of High Concern List

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs - Group 1: Carcinogenic to humans

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs - Group 2B: Possibly carcinogenic to humans

International WHO List of Proposed Occupational Exposure Limit (OEL) Values for Manufactured Nanomaterials (MNMS)

antimony is found on the following regulatory lists

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 4

Australian Inventory of Industrial Chemicals (AIIC)

Chemical Footprint Project - Chemicals of High Concern List

International WHO List of Proposed Occupational Exposure Limit (OEL) Values for Manufactured Nanomaterials (MNMS)

calcium is found on the following regulatory lists

Australia Hazardous Chemical Information System (HCIS) - Hazardous Chemicals

Australian Inventory of Industrial Chemicals (AIIC)

tin fume is found on the following regulatory lists

Australian Inventory of Industrial Chemicals (AIIC)

International WHO List of Proposed Occupational Exposure Limit (OEL) Values for Manufactured Nanomaterials (MNMS)

sulfuric acid is found on the following regulatory lists

Australia Hazardous Chemical Information System (HCIS) - Hazardous Chemicals

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 6

Australian Inventory of Industrial Chemicals (AIIC)

Chemical Footprint Project - Chemicals of High Concern List

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs - Group 1: Carcinogenic to humans

arsenical dust is found on the following regulatory lists

Australia Hazardous Chemical Information System (HCIS) - Hazardous Chemicals

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 4

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 6

Australia Standard for the Uniform Scheduling of Medicines and Poisons (SUSMP) - Schedule 7

Australian Inventory of Industrial Chemicals (AIIC)

FEI Equine Prohibited Substances List - Banned Substances

FEI Equine Prohibited Substances List (EPSL)

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs

International Agency for Research on Cancer (IARC) - Agents Classified by the IARC Monographs - Group 1: Carcinogenic to humans

International WHO List of Proposed Occupational Exposure Limit (OEL) Values for Manufactured Nanomaterials (MNMS)

National Inventory Status

National Inventory	Status
Australia - AIIC / Australia Non-Industrial Use	Yes
Canada - DSL	Yes
Canada - NDSL	No (lead; antimony; calcium; tin fume; sulfuric acid)
China - IECSC	Yes
Europe - EINEC / ELINCS / NLP	Yes
Japan - ENCS	No (lead; antimony; calcium; tin fume; arsenical dust)
Korea - KECI	Yes
New Zealand - NZIoC	Yes
Philippines - PICCS	Yes
USA - TSCA	Yes
Taiwan - TCSI	Yes
Mexico - INSQ	Yes
Vietnam - NCI	Yes
Russia - FBEPH	Yes

Legend:

Yes = All CAS declared ingredients are on the inventory

No = One or more of the CAS listed ingredients are not on the inventory. These ingredients may be exempt or will require registration.

SECTION 16 Other information

Revision Date	22/05/2023
Initial Date	22/05/2023

Other information

Ingredients with multiple cas numbers

Name	CAS No
calcium	7440-70-2, 8047-59-4

Classification of the preparation and its individual components has drawn on official and authoritative sources using available literature references.

The SDS is a Hazard Communication tool and should be used to assist in the Risk Assessment. Many factors determine whether the reported Hazards are Risks in the workplace or other settings. Risks may be determined by reference to Exposures Scenarios. Scale of use, frequency of use and current or available engineering controls must be considered.

Definitions and abbreviations

PC—TWA: Permissible Concentration-Time Weighted Average
PC—STEL: Permissible Concentration-Short Term Exposure Limit
IARC: International Agency for Research on Cancer
ACGIH: American Conference of Governmental Industrial Hygienists
STEL: Short Term Exposure Limit
TEEL: Temporary Emergency Exposure Limit.
IDLH: Immediately Dangerous to Life or Health Concentrations
ES: Exposure Standard
OSF: Odour Safety Factor
NOAEL :No Observed Adverse Effect Level
LOAEL: Lowest Observed Adverse Effect Level
TLV: Threshold Limit Value
LOD: Limit Of Detection
OTV: Odour Threshold Value
BCF: BioConcentration Factors
BEI: Biological Exposure Index
AIIIC: Australian Inventory of Industrial Chemicals
DSL: Domestic Substances List
NDSL: Non-Domestic Substances List
IECSC: Inventory of Existing Chemical Substance in China
EINECS: European INventory of Existing Commercial chemical Substances
ELINCS: European List of Notified Chemical Substances
NLP: No-Longer Polymers
ENCS: Existing and New Chemical Substances Inventory
KECI: Korea Existing Chemicals Inventory
NZIoC: New Zealand Inventory of Chemicals
PICCS: Philippine Inventory of Chemicals and Chemical Substances
TSCA: Toxic Substances Control Act
TCSI: Taiwan Chemical Substance Inventory
INSQ: Inventario Nacional de Sustancias Químicas
NCI: National Chemical Inventory
FBEPH: Russian Register of Potentially Hazardous Chemical and Biological Substances